

ON A PROBLEM OF MORDELL WITH PRIMITIVE ROOTS

CRISTIAN COBELI

ABSTRACT. We consider the sums of the form

$$S = \sum_{x=1}^N \exp((ax + b_1 g_1^x + \cdots + b_r g_r^x)/p),$$

where p is prime and $g_1, \dots, g_r$ are primitive roots $(\bmod p)$. An almost forty years old problem of L. J. Mordell asks to find a nontrivial estimate of S when at least two of the coefficients $b_1, \dots, b_r$ are not divisible by p . Here we obtain a nontrivial bound of the average of these sums when g_1 runs over all primitive roots $(\bmod p)$.

1. INTRODUCTION

Let p be a prime number, $1 \leq N \leq p-1$, r a positive integer and consider the exponential sum

$$S_N(a, \mathbf{b}, \mathbf{g}) := \sum_{x=1}^N e_p(ax + b_1 g_1^x + \cdots + b_r g_r^x), \quad (1.1)$$

where a , and the components of $\mathbf{b} = (b_1, \dots, b_r)$ are integers, $b_1, \dots, b_r$ are not divisible by p and $\mathbf{g} = (g_1, \dots, g_r)$ has components primitive roots modulo p . (We use I. M. Vinogradov's notation $e_p(\alpha) := \exp(2\pi i \alpha/p)$.) In the case $r = 1$, when $p \mid a$ and $p \nmid b$, R. G. Stoneham[5] proved that

$$S_N(b, g) := \sum_{x=1}^N e_p(bg^x) = O(p^{1/2} \log p). \quad (1.2)$$

In a correspondence with D. A. Burgess, L. J. Mordell was informed that both Stoneham and Burgess have found independently several proofs of (1.2). Mordell [3] rediscovered one of the proofs of Burgess and observed that this leads to the following generalization:

$$S_N(a, b, g) := \sum_{x=1}^N e_p(ax + bg^x) < 2p^{1/2} \log p + 2p^{1/2} + 1, \quad (1.3)$$

where $p \nmid ab$. He remarks that his method doesn't seem to apply for the estimate of (1.1) when $r \geq 2$, and the problem remained unsolved till this day. In this paper, fixing all but one of the primitive roots, say $g \in \{g_1, \dots, g_r\}$, we derive a nontrivial bound of $S = S_N(a, \mathbf{b}, \mathbf{g})$ on average over all g primitive roots $(\bmod p)$.

In the following we write shortly $\mathbf{g}^x = (g_1^x, \dots, g_r^x)$, for any integer x and $\mathbf{g} = (g_1, \dots, g_r)$. Also, we use the dot product notation: $\mathbf{b}\mathbf{g}^x = b_1g_1^x + \dots + b_rg_r^x$, where $\mathbf{b} = (b_1, \dots, b_r)$. Let

$$S_N(a, b, \mathbf{b}, g, \mathbf{g}) = \frac{1}{\varphi(p-1)} \sum'_{g \pmod{p}} \sum_{x=1}^N e_p(ax + bg^x + \mathbf{b}\mathbf{g}^x), \quad (1.4)$$

where the prime indicates that the summation is over all g primitive roots $\pmod{p}$.

Theorem 1. *Let p be prime, $1 \leq N \leq p-1$, let $a, b, b_1, \dots, b_r$ be integers not all divisible by p , $\gcd(b, p) = 1$, and let $g, g_1, \dots, g_r$ be primitive roots $\pmod{p}$. Then:*

$$|S_N(a, b, \mathbf{b}, g, \mathbf{g})| \ll p^{\frac{23}{24} + \epsilon}. \quad (1.5)$$

The idea of proof is inspired from the Vinogradov's method and it proved successfully in the estimation of some exponential function analogue of Kloosterman sum, Shparlinski [4].

2. THE COMPLETE INTERVAL CASE

We may assume that $r \geq 1$, since otherwise (1.3) gives a better estimate than (1.5). Taking some fixed primitive root $g_0 \pmod{p}$, then any primitive root $g \pmod{p}$ can be written as $g = g_0^u \pmod{p}$, for some $1 \leq u \leq p-1$ with $\gcd(u, p-1) = 1$. This allows us to replace the sum over g in (1.4) by a sum over $1 \leq u \leq p-1$ with $\gcd(u, p-1) = 1$. Then

$$\begin{aligned} S_N(a, b, \mathbf{b}, g, \mathbf{g}) &= \frac{1}{\varphi(p-1)} \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} \sum_{x=1}^N e_p(ax + bg^{ux} + \mathbf{b}\mathbf{g}^x), \\ &\ll \frac{\Sigma_N}{\varphi(p-1)}, \end{aligned} \quad (2.1)$$

where

$$\Sigma_N = \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} \left| \sum_{x=1}^N e_p(ax + bg^{ux} + \mathbf{b}\mathbf{g}^x) \right|.$$

From now on, in this section we assume that $N = p-1$ and write shortly $\Sigma = \Sigma_{p-1}$. Applying the Cauchy-Schwarz inequality, we have:

$$\begin{aligned}
\Sigma^2 &\leq \varphi(p-1) \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} \left| \sum_{x=1}^{p-1} e_p(ax + bg^{ux} + \mathbf{b}g^x) \right|^2 \\
&= \varphi(p-1) \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} \sum_{x=1}^{p-1} \sum_{y=1}^{p-1} e_p(ax + \mathbf{b}g^x - ay - \mathbf{b}g^y) e_p(bg^{ux} - bg^{uy}) \\
&\leq \varphi(p-1) \sum_{x=1}^{p-1} \sum_{y=1}^{p-1} \left| e_p(a(x-y) + \mathbf{b}g^x - \mathbf{b}g^y) \right| \cdot \left| \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} e_p(b(g^{ux} - g^{uy})) \right|.
\end{aligned}$$

Then, by the Hölder Inequality, we get

$$\begin{aligned}
\Sigma^8 &\leq \varphi(p-1)^4 \left(\sum_{x=1}^{p-1} \sum_{y=1}^{p-1} \left| \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} e_p(b(g^{ux} - g^{uy})) \right| \right)^4 \\
&\leq \varphi(p-1)^4 \left(\sum_{x=1}^{p-1} \sum_{y=1}^{p-1} 1 \right)^3 \sum_{x=1}^{p-1} \sum_{y=1}^{p-1} \left| \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} e_p(b(g^{ux} - g^{uy})) \right|^4.
\end{aligned}$$

Replacing y by xy and then g^x by λ , we have:

$$\begin{aligned}
\Sigma^8 &\leq p^{10} \sum_{x=1}^{p-1} \sum_{y=1}^{p-1} \left| \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} e_p(b(g^{ux} - g^{uxy})) \right|^4 \\
&\leq p^{10} \sum_{\lambda=1}^{p-1} \sum_{y=1}^{p-1} \left| \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} e_p(b(\lambda^u - \lambda^{uy})) \right|^4.
\end{aligned} \tag{2.2}$$

The double sum on y and u can be estimated following the proof of Theorem 8 from Canetti et al [2]. The result is stated in the following lemma:

Lemma 2. *For any integers b , $\gcd(a, b, p) = 1$ and λ primitive root mod p , we have*

$$H_{a,b} = \sum_{y=1}^{p-1} \left| \sum_{\substack{x=1 \\ \gcd(x, p-1)=1}}^{p-1} e_p(a\lambda^x + b\lambda^{xy}) \right|^4 = O(p^{14/3+\epsilon}). \tag{2.3}$$

The estimate (2.3) is a generalization and improvement of Theorem 10 from Canetti, Friedlander, Shparlinski [1].

Proof. Using the properties of the Möbius function and then the Hölder inequality, we have:

$$\begin{aligned}
H_{a,b} &= \sum_{y=1}^{p-1} \left| \sum_{d|p-1} \mu(d) \sum_{\substack{x=1 \\ d|x}}^{p-1} e_p(a\lambda^x + b\lambda^{xy}) \right|^4 \\
&\leq \sum_{y=1}^{p-1} \left(\sum_{d|p-1} \left| \sum_{x=1}^{\frac{p-1}{d}} e_p(a\lambda^{dx} + b\lambda^{dxy}) \right| \right)^4 \\
&\leq \sum_{y=1}^{p-1} \left(\sum_{d|p-1} 1 \right)^3 \sum_{d|p-1} \left| \sum_{x=1}^{\frac{p-1}{d}} e_p(a\lambda^{dx} + b\lambda^{dxy}) \right|^4 \\
&\leq \sigma_0^3(p-1) \sum_{d|p-1} \sum_{y=1}^{p-1} \left| \sum_{x=1}^{\frac{p-1}{d}} e_p(a\lambda^{dx} + b\lambda^{dxy}) \right|^4,
\end{aligned} \tag{2.4}$$

where $\sigma_r(n) = \sum_{d|n} d^r$ is the sum of powers of the divisors of n . For any $d \mid p-1$ we denote $t_d := (p-1)/d$ and $\lambda_d := \lambda^d$. Notice that the multiplicative order of λ^d is t_d . Then the sum over y from the last line above becomes:

$$\begin{aligned}
\sum_{y=1}^{p-1} \left| \sum_{x=1}^{\frac{p-1}{d}} e_p(a\lambda^{dx} + b\lambda^{dxy}) \right|^4 &= d \sum_{y=1}^{t_d} \left| \sum_{x=1}^{t_d} e_p(a\lambda_d^x + b\lambda_d^{xy}) \right|^4 \\
&= d \sum_{y=1}^{t_d} \frac{1}{t_d} \sum_{z=1}^{t_d} \left| \sum_{x=1}^{t_d} e_p(a\lambda_d^{x+z} + b\lambda_d^{(x+z)y}) \right|^4 \\
&= \frac{d}{t_d} \sum_{y=1}^{t_d} \sum_{z=1}^{t_d} \left| \sum_{x=1}^{t_d} e_p(a\lambda_d^z \lambda_d^x + b\lambda_d^{zy} \lambda_d^{xy}) \right|^4 \\
&\leq \frac{d}{t_d} \sum_{y=1}^{t_d} \sum_{\alpha, \beta=0}^{p-1} \left| \sum_{x=1}^{t_d} e_p(\alpha \lambda_d^x + \beta \lambda_d^{xy}) \right|^4,
\end{aligned} \tag{2.5}$$

since for each fixed $y \in \{1, \dots, t_d\}$ the pairs $(a\lambda_d^z, b\lambda_d^{zy})$ with $z \in \{1, \dots, t_d\}$ are distinct modulo p . Next we write explicitly the absolute value in the last term and see that (2.5)

gives

$$\begin{aligned}
& \sum_{y=1}^{p-1} \left| \sum_{x=1}^{\frac{p-1}{d}} e_p(a\lambda^{dx} + b\lambda^{dxy}) \right|^4 \\
& \leq \frac{d}{t_d} \sum_{\alpha, \beta=0}^{p-1} \sum_{y, x_1, x_2, x_3, x_4=1}^{t_d} e_p(\alpha(\lambda_d^{x_1} + \lambda_d^{x_2} - \lambda_d^{x_3} - \lambda_d^{x_4}) + \beta(\lambda_d^{x_1 y} + \lambda_d^{x_2 y} - \lambda_d^{x_3 y} - \lambda_d^{x_4 y})) \\
& \leq \frac{d}{t_d} \sum_{\substack{y, x_1, x_2=1 \\ x_3, x_4=1}}^{t_d} \sum_{\alpha=0}^{p-1} e_p(\alpha(\lambda_d^{x_1} + \lambda_d^{x_2} - \lambda_d^{x_3} - \lambda_d^{x_4})) \sum_{\beta=0}^{p-1} e_p(\beta(\lambda_d^{x_1 y} + \lambda_d^{x_2 y} - \lambda_d^{x_3 y} - \lambda_d^{x_4 y})) \\
& = \frac{d}{t_d} \cdot p^2 \cdot T_d,
\end{aligned} \tag{2.6}$$

where T_d is the number of solutions of the system of congruences:

$$\begin{cases} \lambda_d^{x_1} + \lambda_d^{x_2} \equiv \lambda_d^{x_3} + \lambda_d^{x_4}, \\ \lambda_d^{x_1 y} + \lambda_d^{x_2 y} \equiv \lambda_d^{x_3 y} + \lambda_d^{x_4 y} \end{cases}$$

with $1 \leq x_1, x_2, x_3, x_4, y \leq t_d$. In the proof of Theorem 8 from Canetti et al [2], the last inequality bounds T_d by

$$T_d \ll t_d^{14/3} p^{-1}. \tag{2.7}$$

Then, by (2.4), (2.6) and (2.7), we obtain

$$\begin{aligned}
H_{a,b} & \ll \sigma_0^3(p-1) \sum_{d|p-1} \frac{d}{t_d} \cdot p^2 \cdot t_d^{14/3} p^{-1} \\
& \ll \sigma_0^3(p-1) \sigma_{-\frac{14}{3}}(p-1) \cdot p^{14/3}
\end{aligned}$$

and the lemma follows, since $\sigma_r(n) \ll n^\epsilon$ for any r .

□

By (2.2) and (2.3) we deduce that:

$$\Sigma^8 \ll p^{10} \sum_{\lambda=1}^{p-1} p^{14/3+\epsilon} \ll p^{47/3+\epsilon}.$$

Then making use of the estimate $p/\log \log p \ll \varphi(p-1)$, we obtain

$$\frac{\Sigma}{\varphi(p-1)} \ll p^{23/24+\epsilon}. \tag{2.8}$$

From this estimate together with (2.1), it follows (1.5), so Theorem 1 is proved in the case $N = p - 1$.

3. COMPLETION OF THE PROOF

It remains to show that the size of the incomplete sums is not far from that of the complete ones. Let I be an interval of integers $\subseteq [1, p-1]$ and denote

$$S(I) = \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} \sum_{x \in I} e_p(ax + bg^{ux} + \mathbf{b}g^x). \quad (3.1)$$

In order to estimate the departure of $S(I)$ from $S([1, p-1])$, the following characteristic function of the interval I is suitable:

$$\frac{1}{p} \sum_{y \in I} \sum_{k=1}^p e_p(k(y-x)) = \begin{cases} 1, & \text{if } x \in I; \\ 0, & \text{else.} \end{cases}$$

Then

$$\begin{aligned} S(I) &= \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} \sum_{x \in I} e_p(ax + bg^{ux} + \mathbf{b}g^x) \\ &= \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} \sum_{x=1}^{p-1} e_p(ax + bg^{ux} + \mathbf{b}g^x) \frac{1}{p} \sum_{y \in I} \sum_{k=1}^p e_p(k(y-x)) \\ &= \frac{1}{p} \sum_{k=1}^p \sum_{y \in I} e_p(ky) \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} \sum_{x=1}^{p-1} e_p((a-k)x + bg^{ux} + \mathbf{b}g^x). \end{aligned}$$

In this last form of $S(I)$ we separate the terms with $k = p$ and bound its absolute value to get:

$$\begin{aligned} |S(I)| &\leq \frac{1}{p} \sum_{k=1}^{p-1} \left| \sum_{y \in I} e_p(ky) \right| \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} \left| \sum_{x=1}^{p-1} e_p((a-k)x + bg^{ux} + \mathbf{b}g^x) \right| \\ &\quad + \frac{1}{p} |I| \sum_{\substack{u=1 \\ \gcd(u, p-1)=1}}^{p-1} \left| \sum_{x=1}^{p-1} e_p(ax + bg^{ux} + \mathbf{b}g^x) \right|. \end{aligned} \quad (3.2)$$

Here the sum over y is a geometric progression, that can be evaluated accurately using

$$|e_p(k) - 1| = 2 \left| \sin \left(\frac{k\pi}{p} \right) \right| \geq 4 \left\| \frac{k}{p} \right\|, \quad (3.3)$$

where $\|\cdot\|$ is the distance to the nearest integer, while the sums over u and x are the complete sums bounded by (2.8). Thus, by (3.2), (3.3) and (2.8), we get

$$\begin{aligned}
|S(I)| &\leq \frac{1}{p} \sum_{k=1}^{p-1} \frac{2}{|e_p(k) - 1|} p^{47/24+\epsilon} + \frac{1}{p} |I| p^{47/24+\epsilon} \\
&\leq p^{47/24+\epsilon} \left(\frac{1}{p} \sum_{k=1}^{\frac{p-1}{2}} \frac{1}{2k/p} + 1 \right) \\
&\leq p^{47/24+\epsilon} (3 + \log p) \\
&\leq p^{47/24+\epsilon},
\end{aligned}$$

which concludes the proof of Theorem 1.

Acknowledgement: We thank Ke Gong for his careful reading of an earlier version of this paper.

REFERENCES

- [1] R. Canetti, J. B. Friedlander, and I. E. Shparlinski, *On certain Exponential Sums and the Distribution of Diffie-Hellman Triples*, J. London Math. Soc., **59** (1999), 799–812.
- [2] R. Canetti, J. B. Friedlander, S. Konyagin, M. Larsen, D. Lieman and I. E. Shparlinski, *On the Statistical Properties of Diffie-Hellman Distributions*, Israel J. Math., **120** (2000), 23–46.
- [3] L. J. Mordell, *On the Exponential sum $\sum_{x=1}^X \exp(2\pi i(ax + bg^x)/p)$* , Mathematika, **19** (1972), 84–87.
- [4] I. Shparlinski, *Exponential Function Analogue of Kloosterman Sums*, Rocky Mountain J. Math., 2004, v. **34**, 1497–1502.
- [5] R. G. Stoneham, *On the Uniform ε -Distribution of Residues within the Periods -of Rational Fractions with Applications to Normal Numbers*, Acta Arithmetica, **22** (1972), 371–389.

CRISTIAN COBELI, MATHEMATICS RESEARCH INSTITUTE OF THE ROMANIAN ACADEMY, P.O. BOX 1-764, BUCHAREST, 70700, ROMANIA

E-mail address: cristian.cobeli@imar.ro